

**INDICATIONS:** For use in healthy cattle and calves as an aid in the prevention of disease due to Bovine Virus Diarrhea virus. The vaccine contains BVDV Type 1, with protection demonstrated against BVDV Type 1 challenge.

Contains penicillin and streptomycin as preservatives.

**DIRECTIONS:** Store at 2° to 7° C. Rehydrate with accompanying vial of sterile diluent supplied as a component part of this package. Shake well after rehydration. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.

Cattle should be vaccinated before or when entering feedlots or areas where dairies are located.

Sterilize syringes and needles by boiling in clean water – do not use disinfectants or detergents.

**PRECAUTIONS:** Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

Burn, autoclave, or chemically disinfect this container and all unused contents.

**DOSAGE AND ADMINISTRATION:** Inject 2ml of the combined product intramuscularly (preferably in the neck region).

Do not administer to pregnant cows or calves nursing pregnant cows.

Yearly booster injections are recommended for breeding stock. Be sure that breeding stock are vaccinated on a regular basis when cows are open.

Maternal antibodies in young calves may interfere with development of an active immunity. If emergency conditions require vaccination these calves should be revaccinated one month after weaning.

## BOVINE VIRUS DIARRHEA VACCINE


BVD

Modified Live Virus

Cat. #19402	20ml	10 Dose
Cat. #19404	100ml	50 Dose



**FOR VETERINARY USE ONLY**



[www.coloradoserum.com](http://www.coloradoserum.com)