

**INDICATIONS:** For use in healthy equines as an aid in the prevention of Eastern and Western types of Encephalomyelitis and tetanus.

Formalin inactivated adjuvanted equine encephalomyelitis vaccine in combination with concentrated tetanus toxoid. Contains thimerosal, penicillin, and streptomycin as preservatives.

**DIRECTIONS:** Store at 2° to 7° C. Do not freeze. Do not vaccinate within 21 days before slaughter.

Shake well before use. Use entire contents when first opened.

**PRECAUTIONS:** Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

**DOSAGE AND ADMINISTRATION:** Inject two 2ml doses intramuscularly with an interval of 3 weeks between injections. Revaccinate annually using a single dose and whenever an epidemic situation develops and exposure is likely.

Encephalomyelitis is spread by biting insects, so animals should be vaccinated prior to the time insects become prevalent.

## ENCEPHALOMYELITIS VACCINE

Killed Virus

### EASTERN & WESTERN TETANUS TOXOID

Cat. #18211 10 X 2ml 10 X 1 Dose

Cat. #18215 20ml 10 Dose



**FOR VETERINARY USE ONLY**



[www.coloradoserum.com](http://www.coloradoserum.com)