

## ESSENTIAL 2+P

Clostridium Chauvoei-Septicum  
Mannheimia Haemolytica-Pasteurella-  
Multocida Bacterin  
Killed Bacterin

Cat. #11352 50ml 10 Dose

Cat. #11354 250ml 50 Dose



### FOR VETERINARY USE ONLY



**INDICATIONS:** For use in healthy cattle, sheep and goats as an aid in the prevention of Blackleg, Malignant Edema, and Pasteurellosis caused by the micro-organisms named.

Product is aluminum hydroxide adsorbed. Contains thimerosal as a preservative.

**DIRECTIONS:** Store at 2° to 7° C. Do not freeze. Shake well before use. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.

**PRECAUTIONS:** Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

### DOSAGE AND ADMINISTRATION:

**Cattle:** Inject 5ml subcutaneously. Calves vaccinated under three months of age should be revaccinated at weaning or 4 to 6 months of age.

**Sheep and Goats:** Inject 3ml subcutaneously. Revaccinate with Mannheimia Haemolytica-Pasteurella Multocida Bacterin in 2 to 4 weeks.

