

INDICATIONS: For use in healthy cattle, sheep and goats as an aid in the prevention of enterotoxemia caused by *Clostridium perfringens* Types B, C, and D. *Clostridium perfringens* Type B is not a significant problem in North America.

For use in healthy swine as an aid in the prevention of enterotoxemia caused by *Clostridium perfringens* Type C.

For the long-term protection against tetanus.

Contains thimerosal as a preservative.

DIRECTIONS: Store at 2° to 7° C. Do not freeze. Shake well before use. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.

Vaccinate sufficiently in advance of feeding concentrated rations to provide a minimum of 2 weeks after second dose for adequate immunity to develop.

Tetanus toxoid requires 3 to 4 weeks to establish effective protection that will last several months. Booster injections should be made annually, or, in event of injury, regardless of interval.

PRECAUTIONS: Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

DOSAGE AND ADMINISTRATION: Inject 2 ml subcutaneously or intramuscularly. Repeat full dose in 3 to 4 weeks.

ESSENTIAL 3+T
Clostridium Perfringens Types C&D
Tetanus Toxoid
Detoxified Toxin

Cat. #11302	20ml	10 Dose
Cat. #11304	100ml	50 Dose
Cat. #11305	250ml	125 Dose



FOR VETERINARY USE ONLY

