

INDICATIONS: For use in healthy calves and non-pregnant cattle as an aid in the prevention of infectious Bovine Rhinotracheitis and Leptospirosis caused by the serotypes included in the bacterin.

Contains thimerosal, penicillin, and streptomycin as preservatives.

DIRECTIONS: Store at 2° to 7° C.

Rehydrate the vaccine just prior to use. To rehydrate, withdraw the entire contents of the accompanying Lepto-5 Bacterin vial with a sterile syringe and needle, without removing the stopper, and transfer it to the vial of dried vaccine. Shake well after rehydration. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.

Do not administer to pregnant cows or calves nursing pregnant cows. Do not administer to calves less than 4 weeks of age. Maternal antibodies in young calves may interfere with development of active immunity. If necessary to vaccinate young calves, revaccinate these animals at 6 months of age.

Cattle should be vaccinated before or when entering feedlots or areas where dairies are located.

Yearly booster injections are recommended for breeding stock. Be sure that breeding stock are vaccinated on a regular basis when cows are open.

Sterilize syringes and needles by boiling in clean water – do not use disinfectants or detergents.

PRECAUTIONS: Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

Burn, autoclave, or chemically disinfect this container and all unused contents.

DOSAGE AND ADMINISTRATION: Inject 2ml of the combined product intramuscularly (preferably in the neck region).



PRE-BREED 6 IBR-LEPTO 5

Modified Live Virus/Killed Bacterin

Cat. #19532 20ml 10 Dose
Cat. #19534 100ml 50 Dose



FOR VETERINARY USE ONLY



www.coloradoserum.com