



RESPIRA-3 IBR-BVD-PI₃ Bovine Rhinotracheitis-Virus Diarrhea-Parainfluenza ₃ Vaccine Modified Live Virus		
Cat. #19452	20ml	10 Dose
Cat. #19454	100ml	50 Dose
		

FOR VETERINARY USE ONLY



INDICATIONS: For use in healthy cattle and calves as an aid in the prevention of disease caused by bovine rhinotracheitis virus, bovine virus diarrhoea virus (BVDV) and Parainfluenza₃ virus. The vaccine contains BVDV Type 1, with protection demonstrated against BVDV Type 1 challenge.

Contains penicillin and streptomycin as preservatives.

DIRECTIONS: Store at 2° to 7° C. Rehydrate with accompanying vial of sterile diluent supplied as a component part of this package. Shake well after rehydration. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.

Do not administer to pregnant cows or calves nursing pregnant cows. Do not administer to calves less than 4 weeks of age.

Cattle should be vaccinated before or when entering feedlots or areas where dairies are located.

Yearly booster injections are recommended for breeding stock. Be sure that breeding stock are vaccinated on a regular basis when cows are open.

Sterilize syringes and needles by boiling in clean water – do not use disinfectants or detergents.

PRECAUTIONS: Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

Burn, autoclave, or chemically disinfect this container and all unused contents.

DOSAGE AND ADMINISTRATION: Inject 2ml of the combined product intramuscularly (preferably in the neck region).

Maternal antibodies in young calves may interfere with development of active immunity. If necessary to vaccinate young calves, revaccinate these animals at 6 months of age.