


TETANUS TOXOID
Concentrated, Adjuvanted

Detoxified Toxin

Cat. #11411 10 X 1ml 10 X 1 Dose
Cat. #11415 10ml 10 Dose



FOR VETERINARY USE ONLY



INDICATIONS: For use as an aid in the prevention of tetanus. Contains thimerosal as a preservative.

DIRECTIONS: Store at 2° to 7° C. Do not freeze. Shake well before use. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter.

Vaccination with Tetanus Toxoid is recommended for healthy domestic animals, not infected with tetanus, to establish an active immunity for prevention against disease. Protective antibody levels usually occur about two weeks after the second injection of the primary immunization series.

In contrast, administration of Tetanus Antitoxin is recommended for immediate, emergency, passive treatment of exposed animals with an unknown vaccination history or with signs of tetanus infection. Refer to the Tetanus Antitoxin product circular for full information and consult with a veterinarian.

PRECAUTIONS: Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

DOSAGE AND ADMINISTRATION: For primary immunization, two doses should be administered subcutaneously or intramuscularly approximately 30 days apart. Use intramuscularly for horses, as local reactions are more likely to occur if injected subcutaneously.

Cattle and Horses: 1 ml dose
Sheep, Goats, and Swine: 0.5 ml dose

Revaccinate annually using a single dose.

